



Coltox

Inactivated vaccine against Necrotic Enteritis and Neonatal E. Coli Diarrhea in injectable suspension



Composition per dose (2 ml)

Inactivated *E. coli* strains: $\geq 2 \times 10^9$ CFU
for each strain with the following fimbrial
antigens: K88(F4), K99(F5), P987(F6) and F41

Clostridium perfringens

C and D: ≥ 10 I.U. of antitoxin β per ml of rabbit serum
 ≥ 5 I.U. of antitoxin ϵ per ml of rabbit serum

Indications

Prevention of enterotoxemia and diarrhoea in piglets caused by different serotypes of enterotoxigenic *Escherichia Coli* and *Clostridium perfringens* types C and D.

Administration Route

Deep intramuscular injection.

Dosage and administration

Sows and gilts: 2 ml.

Piglets:

First dose: 0,5 ml.

Second dose: 1 ml .

Pregnant sows: administer one dose 6-7 weeks prior to farrowing, and a second dose 4 weeks before farrowing. Then administer a single dose at every pregnancy, 4 weeks before farrowing.

Piglets: administer 0,5 ml at 10 days of age and a second dose of 1 ml at weaning.

Shelf-life period

Two years.

Withdrawal period

Zero days.

Storage

Store and transport this product refrigerated (between 2°C and 8°C). Do not freeze.

Supply

Vials of 50 ml (25 doses) and 100 ml (50 doses).

LICENSE NUMBER: B-7025-015 SAGARPA (MEXICO)

syva