



Agalax Tres

Inactivated vaccine against Contagious Agalactia in injectable suspension



Composition per dose (2 ml)

<i>Mycoplasma agalactiae</i>	≥ 1:20 *
<i>Mycoplasma capricolum</i>	≥ 1:20 *
<i>Mycoplasma mycoides</i> subespecie <i>mycoides</i>	≥ 1:20 *

* Antibody titre obtained by the Growth Inhibition Technique after vaccinating lambs with one dose of the vaccine.

Indications

Sheep and goats: active immunization of sheep and goats against Contagious Agalactia caused by *Mycoplasma agalactiae*, *Mycoplasma capricolum*, *Mycoplasma mycoides* subsp. *mycoides*.

Administration Route

Subcutaneous injection.

Dosage and administration

Dose: 2 ml per animal.

In flocks free of the disease: administer one dose to all male and female breeders, and to all reposition animals from their 2nd or 3rd month of life onwards.

In infected flocks or if the risk of infection is high: apply a booster injection 2-4 weeks after the first administration.

In general a booster injection is recommended every 4 to 6 months depending on the epidemiological situation.

Shelf-life period

Two years.

Withdrawal period

Zero days.

Storage

Store and transport this product refrigerated (between 2°C and 8°C). Do not freeze.

Supply

Vials of 100ml. (50 doses) and 250 ml. (125 doses).

LICENSE NUMBER: 7929 NAL (AEM SPAIN)