



# Facilpart

## Injectable Oxytocin



### Pharmaceutical Form

Injectable solution.

### Composition per ml

Synthetic Oxytocin

10 I.U.

### Indications

**In general and for all species:** induction to parturition, uterus atony, uterus involution, metritis, retained placenta, lactation induction after parturition, milk ejection, agalaxia in sows. Also as coadjuvant therapy for the antibiotic treatment of acute and chronic mastitis, in order to facilitate the expulsion of residues and facilitating the drainage.

### Administration Route

Intramuscular, intravenous or subcutaneous injection.

### Dosage and administration

Obstetrics (intravenous, intramuscular or subcutaneous injection):

**Cows:** 7,5-10 ml per cow.

**Mares:** 7,5-15 ml per mare.

**Sows, sheep and goats:** 3-5 ml per animal.

**Bitch:** 0,5-2,5 ml per bitch.

**Female cats:** 0,5-1 ml per cat.

Milk ejection (preferably intravenously):

**Cows and mares:** 1-2 ml per animal.

**Sows, sheep and goats:** 0,5-2 ml per animal.

**Bitch:** 0,2-1 ml per bitch.

**Female cats:** 0,1-1 ml per cat.

The administration can be repeated every 30 minutes if the veterinary surgeon considers it necessary.

### Withdrawal period

Meat: 2 days.

Milk: 2 days.

### Shelf-life period

2 years.

### Storage

In a cool, dark place.

### Supply

Vials of 10, 100 and 250 ml.

LICENSE NUMBER: 0528-ESP (AEM SPAIN)