



Flumesyva líquido 20%

Flumequine 20% Oral Solution



Pharmaceutical Form

Oral solution.

Composition per ml

Flumequine.

200 mg.

Indications

Treatment of germs sensitive to flumequine, which causes infections such as:

Poultry and turkey: colibacillosis, salmonellosis and pasteurellosis.

Pigs: colibacillosis, enteritis, gastroenteritis, neonatal diseases.

Lambs: colibacillosis, septicaemia, pasteurellosis and pneumonia.

Calves: bronchopneumonia, colibacillosis enteritis, neonatal diseases and salmonellosis.

Administration Route

Orally, by addition to drinking water.

Dosage and administration

Poultry: 1 ml per 8-16 kg body weight during 5 days, equivalent to 0.5 ml per litre and day of drinking water.

Adult pig: 0,30 ml per 10 kg of body weight, every 12 hours, during 3-5 days.

Piglet: 0,60 ml per 10 kg of body weight, every 12 hours, during 3-5 days.

Lambs: 0,30 ml per 10 kg of body weight, every 12 hours, during 4-6 days.

Calves: 0,25-0,50 ml per 10 kg of body weight, every 12 hours, during 5 days.

Withdrawal period

Meat: 10 days.

Eggs: do not use in laying hens.

Shelf-life period

3 years.

Storage

Store in a cool, dark place.

Supply

Bottle of 1 litre.

LICENSE NUMBER: 0571-ESP (AEM SPAIN)