



Flumesyva polvo 10%

Flumequine 10%, Water Soluble Powder



Pharmaceutical Form

Water soluble powder.

Composition per gram

Flumequine 100 mg.

Indications

Treatment of microorganisms sensitive to flumequine, which cause infections such as:

Poultry and turkey: colibacillosis, salmonellosis and pasteurellosis.

Pigs: colibacillosis, enteritis, gastro-enteritis, neonatal diseases.

Calves: bronchopneumonia, colibacillosis, bacterial enteritis, neonatal diseases and salmonellosis.

Lambs: colibacillosis, septicaemia, pasteurellosis and pneumonia.

Fish: furunculosis in salmonidae.

Administration Route

Oral, by addition to feed or to drinking water.

Dosage and administration

Poultry and turkey: 1 g per 4-8 kg of body weight and day, during 3-5 days.

Adult swine: 3 g per 50 kg of body weight, every 12 hours, during 3-5 days.

Piglets: 3 g per 25 kg of body weight, every 12 hours, during 3-5 days.

Lambs: 0,60 g per 10 kg of body weight, every 12 hours, during 4-6 days.

Calves: 5 g per 50 to 100 kg of body weight, every 12 hours, during 5 days.

Fish: 0,12 g per kg of body weight and day.

Withdrawal period

Pigs and poultry: 2 days.

Turkey, lambs, calves and fish: 10 days.

Eggs: do not use in laying hens.

Shelf-life period: 3 years.

Storage: in a cool, dry and dark place.

Supply

Box of 1 kg (10 bags of 100 g.).

Box of 5 kg (5 bags of 1 kg).

LICENSE NUMBER: 0570-ESP (AEM SPAIN)

syva