



Gentayet

Injectable Gentamycin



Pharmaceutical Form

Injectable solution.

Composition per ml

Gentamycin (sulphate)

40 mg.

Indications

Cattle: coli-mastitis, colibacillosis and salmonellosis in calves, genitourinary diseases, respiratory tract infections.

Horses: colibacillosis and salmonellosis in foals, genitourinary infections and respiratory tract infections.

Piglets: neonatal colibacillosis.

Dogs: genitourinary infections, respiratory infections, infections of the skin and soft tissues, bacterial gastroenteritis, bacteremia and septicaemia.

Cats: genitourinary infections, respiratory infections, infections of the skin and soft tissues, bacterial gastroenteritis.

Administration Route

Deep intramuscular or slow intravenous injection.

Dosage and administration

Cattle: 1 ml per 8-10 kg body weight, every 12 hours, for three days.

Horses: 1 ml per 10-20 kg body weight, every 8 hours, for three days.

Piglets: a single dose of 0,125 ml per animal.

Dogs and Cats: 1 ml per 8 kg body weight, every 12 hours, for 3-4 days.

Withdrawal period

Cattle: 60 days.

Swine: 40 days.

Milk: 3 days.

Do not administer in horses whose meat is for human consumption.

Shelf-life period

3 years.

Storage

Store in a cool, dark place.

Supply

Vials of 100 ml and 250 ml.