



Sylvacal 20

Oral Calcium + Magnesium + Vitamin D3



Pharmaceutical Form

Oral suspension.

Composition per 500 ml

Dihydrate calcium chloride	203 g. (55 g. of calcium).
Hexahydrate magnesium chloride	8 g. (1 g. of magnesium).
Cholecalciferol (vitamin D ₃)	100.000 I.U.

Indications

Cattle:

SYVACAL 20 is indicated in the prophylaxis and treatment of hypocalcaemia in general. It is also used as a support to intravenous calcium therapy and for the prophylaxis of relapses.

Its use is highly recommended just before and after calving, and at lactation initial stages.

Administration Route

Oral administration, as deep as possible inside the oral cavity.

Dosage and administration

Parturition and lactation preparation: administer in total 4 bottles per cow. One day before calving, another immediately after calving, and again 12 and 24 hours later.

As a dietetic complement to the injectable calcium: 2 bottles per cow, administered 12 and 24 hours after the treatment with injectable calcium.

Prophylaxis and therapy of hypocalcaemia: 2-4 bottles per cow at intervals of 12 hours.

Withdrawal period

Not applicable.

Shelf-life period

3 years.

Storage

No special conditions are applicable.

Supply

Bottle of 500 ml.

LICENSE NUMBER: LICENSED AS FEED ADDITIVE