



Syvayesky acuoso

Liophilized Live Attenuated vaccine against Pseudorabies (Aujeszky's Disease) with aqueous adjuvant



Composition per dose (2 ml)

Aujeszky Disease Virus, Bartha strain K61 gE⁻ $\geq 10^6$ TCID₅₀
Aqueous adjuvant

Indications

Active immunization of pigs against Aujeszky's Disease.

Administration

Deep intramuscular injection or intra-nasal administration.

Dosage and administration

Dose: 2 ml per pig.

Piglets and fatteners: first vaccination at 10 to 12 weeks of age.

Second vaccination 3 to 4 weeks later. Third vaccination at 6 month of age. Revaccination every four months.

Gilts: first vaccination at 10 to 12 weeks of age. Second vaccination 3 to 4 weeks later. Third vaccination at 21- to 24 weeks of age.

Sows and boars: vaccinate every four months; that is three vaccinations per year as herd vaccination.

Shelf-life period

One year.

Withdrawal period

Zero days.

Storage

Store and transport this product refrigerated (between 2°C and 8°C). Do not freeze.

Supply

50 doses box (1 vial of 50 doses).

500 doses box (10 vials of 50 doses).

1000 doses box (20 vials of 50 doses).

1000 doses box (10 vials of 100 doses).

LICENSE NUMBER: 10863 NAL (AEM SPAIN)