



# Syvazul 1+8

Inactivated vaccine against the serotype 1+8 of the Bluetongue Disease in injectable suspension



## Composition per ml

Inactivated bluetongue virus, strain ALG2006/01E1 (serotype 1)  $\geq 10^{6.9}$  TCID<sub>50</sub>\*  
Inactivated bluetongue virus, strain BEL2006/01 (serotype 8)  $\geq 10^{6.7}$  TCID<sub>50</sub>\*  
\* Determined before inactivation.

## Indications

Active immunization of sheep and cattle for the prevention of viraemia caused by serotypes 1 and 8 of the Bluetongue Disease virus.

## Administration Route

**Sheep:** subcutaneous injection.

**Cattle:** intramuscular injection.

## Dosage and administration

Dosage: 2 ml for sheep and 4 ml for cattle.

### Initial vaccination:

Administer two doses with an interval of 3 weeks in between.

Minimum vaccination age: Sheep: 3 months.

Cattle: 2 months.

### Booster:

Administer one dose before every risk season.

When used within the frame of an official control program, the vaccination schedule must follow the plan stated by the relevant sanitary authority.

## Shelf-life period

One year.

## Withdrawal period

Zero days.

## Storage

Store and transport under refrigerated conditions (between +2° C and + 8° C) and protected from light. Do not freeze.

## Supply

Vials of 80 and 200 ml.

LICENSE NUMBER: 2056 ESP (AEM SPAIN)