



Synparv-MRL

Inactivated vaccine against Porcine Parvovirus; Leptospirosis and Swine Erysipelas in injectable emulsion



Composition per dose (2 ml)

<i>Porcine Parvovirus</i>	$\geq 10^{6,9}$ TCID ₅₀
<i>Erysipelothrix rhusiopathiae</i> (serotype 1)	$\geq 10^{10}$ CFU
<i>Erysipelothrix rhusiopathiae</i> (serotype 2)	$\geq 10^{10}$ CFU
<i>Leptospira canicola</i>	$\geq 5 \times 10^8$ CFU
<i>Leptospira grippotyphosa</i>	$\geq 5 \times 10^8$ CFU
<i>Leptospira pomona</i>	$\geq 5 \times 10^8$ CFU
<i>Leptospira icterohaemorrhagiae</i>	$\geq 5 \times 10^8$ CFU
<i>Leptospira bratislava</i>	$\geq 5 \times 10^8$ CFU

Indications

Active immunisation against Porcine Parvovirus infection, Leptospirosis and Swine Erysipelas in healthy pigs.

Administration Route

Intramuscular injection.

Dosage and administration

Dose: 2 ml per pig.

In general the following vaccination program is recommended:

Gilts: first doses at 6-7 months of age. Booster dose 3-4 weeks later.

Sows: vaccination 2-4 weeks before AI or mating.

Boars: first doses at 6-7 months of age. Administer a booster dose 3-4 weeks later. Vaccinate again every 6 months.

Shelf-life period

Two years.

Withdrawal period

Zero days.

Storage

Store and transport this product refrigerated (between 2°C and 8°C). Do not freeze.

Supply

Vials of 10 ml (5 doses) and 100 ml (50 doses).

LICENSE NUMBER: B-7025-005 (SAGARPA MEXICO)