



Actionis

Ceftiofur injectable suspension



Pharmaceutical Form

Injectable suspension.

Composition per ml

Ceftiofur (as hydrochloride) 50.0 mg

Indications

Infections associated with bacteria sensitive to ceftiofur:

Pigs: For the treatment of bacterial respiratory disease associated with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *Streptococcus suis*.

Cattle:

For the treatment of bacterial respiratory disease associated with *Pasteurella haemolytica* (*Mannheimia* spp.), *Pasteurella multocida* and *Haemophilus somnus*.

For the treatment of acute interdigital necrobacillosis (panaritium, foot rot), associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus* (*Porphyromonas asaccharolytica*).

For treatment of the bacterial component of acute post-partum (puerperal) metritis within 10 days after calving associated with *Escherichia coli*, *Arcanobacterium pyogenes* and *Fusobacterium necrophorum*, sensitive to ceftiofur.

Administration Route

Swine: intramuscular injection. **Cattle:** subcutaneous injection.

Dosage and administration

Pigs: 1 ml/16 kg bw for 3 days.

Cattle: Respiratory disease: 1 ml/50 kg bw for 3 to 5 days.

Acute interdigital necrobacillosis: 1 ml/50 kg bw at each injection for 3 days.

Withdrawal period

Swine: Meat: 6 days.

Cattle: Meat: 6 days, Milk: Zero days.

Shelf life period

2 years.

Storage

Store in a cool and dark place.

Supply

Vials of 100 and 250 ml.

LICENSE NUMBER: 2273-ESP