



Florfenis®

Antibacterial in injectable solution



Pharmaceutical Form

Injectable solution.

Composition per ml

Florfenicol

300 mg

Indications

Disease caused by bacteria sensitive to Florfenicol.

Cattle: treatment of respiratory infections caused by *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*.

Swine: treatment of acute outbreaks of swine respiratory disease caused by *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Administration Route

Cattle: intramuscular or subcutaneous. **Swine:** intramuscular.

Dosage and administration

Cattle:

Intramuscular administration: Administer 1 ml of FLORFENIS per 15 kg of body. Repeat the treatment 48 hours later.

Subcutaneous administration: Administer 2 ml of FLORFENIS per 15 kg of body weight only once.

Do not administer more than 10 ml in one single injection site. The product shall be administered only in the neck.

Swine:

Intramuscular administration: 1 ml of FLORFENIS / 20 kg of body weight in the neck musculature. Repeat the treatment 48 later.

Do not administer more than 3 ml in one single injection site.

Withdrawal period

Cattle: Meat: 30 days after intramuscular injection. 44 days after subcutaneous injection. Milk: Do not use.

Swine: Meat: 18 days.

Shelf-life period

2 years.

Storage

Keep the vial in the outer carton in order to protect from light. Do not freeze.

Supply

Vials of 100 and 250 ml.

LICENSE NUMBER: 2.542-ESP (AEM SPAIN)