



Respibov

Inactivated Polyvalent vaccine against Respiratory and Reproductive Diseases in Cattle in injectable suspension



Composition per dose (4 ml)

Virus of the Infectious Bovine Rhinotracheitis (IBR)	≥ 1:8 SN *
Virus of Bovine Viral Diarrhoea (BVD)	≥ 1:32 SN *
Virus of Bovine Parainfluenza type 3 (PI-3)	≥ 1:64 SN *
<i>Mannheimia haemolytica</i>	≥ 1:80 ELISA **
<i>Pasteurella multocida</i>	≥ 1:80 ELISA **

* Minimum titre of Serum-Neutralization antibodies obtained after the inoculation in bovine.

** Minimum titre of ELISA antibodies obtained after the inoculation in bovine.

Indications

Cattle: prevention of the Infectious Bovine Rhinotracheitis, Bovine Viral Diarrhoea, Parainfluenza Virus and pneumonia processes in cattle originated by *Mannheimia haemolytica* and *Pasteurella multocida*.

Administration Route

Intramuscular or subcutaneous injection.

Dosage and administration

Dose: 4 ml per animal.

Calves: administer two doses 3 weeks apart. Calves that have been vaccinated before weaning shall be vaccinated again after weaning.

Cows: administer two doses 3 weeks apart. Revaccinate once a year. Cows can be vaccinated in any pregnancy stage.

Shelf-life period

Two years.

Withdrawal period

Zero days.

Storage

Store and transport this product refrigerated (between 2°C and 8°C). Do not freeze.

Supply

Blister with 4 vials of 4 ml (4 x 1 doses)

Vials of 100 ml (25 doses).

LICENSE NUMBER: 2862 ESP (AEM SPAIN)