



Syvaquinol-100 injectable

Enrofloxacin 10% Injectable Solution



Pharmaceutical Form

Injectable solution.

Composition per ml

Enrofloxacin

100 mg

Indications

Pigs: respiratory infections caused by *Pasteurella multocida*, *Mycoplasma spp.* and *Actinobacillus pleuropneumoniae*.

Treatment of Postpartum Dysgalactia Syndrome (PPDS) in sows.

Digestive infections and septicaemia caused by *E.coli*.

Cattle: Respiratory infections caused by *Mannheimia haemolytica*, *Pasteurella multocida* and *Mycoplasmas spp.*

Digestive infections and septicaemia caused by *E.coli*.

Acute arthritis caused by *Mycoplasma spp.*

Administration Route

Pigs: Intramuscular administration

Cattle: Subcutaneous or intravenous administration.

Dosage and administration

Pigs: administer intramuscularly 0,5 ml per 20 kg body weight and day for 3 days.

For the treatment septicemia or digestive infections caused by *E. coli* administer 1 ml per 20 kg body weight and day for 3 days.

Do not administer more than 3 ml in the same injection site.

Cattle: administer subcutaneously or slowly intravenously 1ml per 10 kg body weight and day, for 3-5 days.

Do not administer subcutaneously more than 10 ml per injection site.

Withdrawal period

Cattle:

Meat: 12 days for intramuscular or subcutaneous administration.

5 days for intravenous administration.

Milk: 4 days for subcutaneous administration. 3 days for intravenous administration.

Pig: Meat: 13 days.

Shelf-life period

3 years.

Storage

Store in a cool, dark place.

Supply

Vials of 100 ml and 250 ml.