



Aquamicin polvo

Antibacterial therapy in water soluble powder



Pharmaceutical Form

Water soluble powder.

Composition per gram

Neomycin (sulphate)

490 mg

Indications

Calves: colibacillosis and salmonellosis.

Piglets: colibacillosis, salmonellosis, vibronic dysentery and oedemas disease.

Foals: bacterial infections of the gastrointestinal tract.

Dogs: bacterial infections of the gastrointestinal tract.

Poultry: unspecific diarrhoeas and salmonelosis.

Administration Route

Oral by addition to drinking water, milk, milk- replacer or directly into the mouth.

Dosage and administration

Calves: 1 g per 28-70 kg of B.W. and day during 3 to 5 days.

Piglets: 1 g per 28 kg of B.W. and day during 3 to 5 days.

Foals: 1 g per 70-140 kg of B.W. and day during 3 to 5 days.

Poultry: 0,032% in feed or drinking water during 3 to 5 days.

Dogs: 0,5 g per 14-35 kg of B.W. and day during 3 to 5 days.

Withdrawal period

Meat: Calves: 30 days.

Pigs: 20 days.

Poultry: 5 days.

Do not administer in horses whose meat is for human consumption.

Shelf-life period

3 years.

Storage

Store in a cool, dark place.

Supply

Bags of 1 and 5 kg.

LICENSE NUMBER: 1242-ESP (AEM SPAIN)